

REFINEMENTS OF SOME CLASSICAL INEQUALITIES INVOLVING SINC AND HYPERBOLIC SINC FUNCTIONS

YOGESH J. BAGUL, SUMEDH B. THOOL , CHRISTOPHE CHESNEAU,
RAMKRISHNA M. DHAIGUDE

Abstract. Several bounds of trigonometric-exponential and hyperbolic-exponential type for sinc and hyperbolic sinc functions are presented. In an attempt to generalize the results, some known inequalities are sharpened and extended. Hyperbolic versions are also established, along with extensions.

1. Introduction

Consider the sinc function defined by $\text{sinc } x = (\sin x)/x$, for $x \neq 0$ and $\text{sinc } x = 1$, for $x = 0$. A hyperbolic sinc function is defined similarly. Let us now cite some inequalities for sinc and hyperbolic sinc functions pertaining to the main results of this paper. First, the classical inequalities

$$(1.1) \quad \cos\left(\frac{x}{\sqrt{3}}\right) < \frac{\sin x}{x} < \cos\left(\frac{2}{\pi} \arccos\left(\frac{2}{\pi}\right) \cdot x\right), \quad 0 < x < \frac{\pi}{2},$$

were established by K.S.K. Iyengar, B.S. Madhava Rao and T.S. Nanjundiah in a little-known paper [9]. See also [14]. Recently, J. Sándor ([18]) offered

Received: 25.03.2022. Accepted: 02.11.2022. Published online: 23.11.2022.

(2020) Mathematics Subject Classification: 26D05, 26D07, 33B10.

Key words and phrases: trigonometric-exponential, hyperbolic-exponential, Mitrinović-Adamović inequality, Lazarević inequality, Iyengar-Madhava Rao-Nanjundiah inequality.

©2022 The Author(s).

This is an Open Access article distributed under the terms of the Creative Commons Attribution License CC BY (<http://creativecommons.org/licenses/by/4.0/>).

a new proof to the left inequality of (1.1) and proved its hyperbolic counterpart as follows:

$$(1.2) \quad \cosh\left(\frac{x}{\sqrt{3}}\right) < \frac{\sinh x}{x}, \quad x > 0.$$

R. Klén, M. Visuri, and M. Vuorinen ([10]) found the following inequalities

$$(1.3) \quad \left[\cos\left(\frac{x}{2}\right)\right]^2 < \frac{\sin x}{x} < \left[\cos\left(\frac{x}{3}\right)\right]^3, \quad 0 < x < \frac{\pi}{2}.$$

Y. Lv, G. Wang, and Y. Chu ([11]) obtained the following:

$$(1.4) \quad \left[\cos\left(\frac{x}{2}\right)\right]^{4/3} < \frac{\sin x}{x} < \left[\cos\left(\frac{x}{2}\right)\right]^a, \quad 0 < x < \frac{\pi}{2},$$

where $a = (\ln(\pi/2))/(\ln \sqrt{2}) \approx 1.30299$.

The left inequality of (1.4) is sharper than the corresponding left inequality of (1.3), whereas the right inequality of (1.3) is better than that of (1.4). The analogous inequality to (1.4) is the following one:

$$(1.5) \quad \left[\cosh\left(\frac{x}{2}\right)\right]^{4/3} < \frac{\sinh x}{x} < \cosh^3 x, \quad x > 0,$$

which can be seen in [15, 16, 20]. Exponential-type bounds for sinc and hyperbolic sinc functions were obtained by Chesneau and Bagul in [6]. They are given below. We have

$$(1.6) \quad e^{\gamma x^2} < \frac{\sin x}{x} < e^{-x^2/6}, \quad 0 < x < \frac{\pi}{2},$$

where $\gamma = 4 \ln(2/\pi)/\pi^2$, and

$$(1.7) \quad e^{\lambda x^2} < \frac{\sinh x}{x} < e^{x^2/6}, \quad 0 < x < r,$$

where $r > 0$ and $\lambda = \ln[(\sinh r)/r]/r^2$.

For different refinements, generalizations, and recent developments regarding inequalities involving the sinc and hyperbolic sinc functions, we refer the reader to [2–5, 7, 12, 13, 17, 21–26]. This article aims to present new generalized bounds for sinc and hyperbolic sinc functions. Our bounds are trigonometric-exponential and hyperbolic-exponential in nature and they refine some existing bounds in the literature.

We consider the following plan: Section 2 presents some preliminaries and lemmas. The main results are given in Section 3. Section 4 ends the article with some particular cases and discussions.

2. Preliminaries and lemmas

The following power series expansions involving Bernoulli numbers can be found in [8, 1.411]:

$$(2.1) \quad \tan x = \sum_{n=1}^{\infty} \frac{2^{2n}(2^{2n} - 1)}{(2n)!} |B_{2n}| x^{2n-1}, \quad |x| < \frac{\pi}{2},$$

$$(2.2) \quad \cot x = \frac{1}{x} - \sum_{n=1}^{\infty} \frac{2^{2n}}{(2n)!} |B_{2n}| x^{2n-1}, \quad |x| < \pi,$$

and

$$(2.3) \quad \tanh x = \sum_{n=1}^{\infty} \frac{2^{2n}(2^{2n} - 1)}{(2n)!} B_{2n} x^{2n-1}, \quad |x| < \frac{\pi}{2},$$

where B_{2n} are the even indexed Bernoulli numbers.

We will also use the following l'Hôpital's rule of monotonicity.

LEMMA 1 (l'Hôpital's rule of monotonicity [1]). *Let $f, g: [a, b] \rightarrow \mathbb{R}$ be two continuous functions which are differentiable on (a, b) and $g' \neq 0$ on (a, b) . If f'/g' is increasing (or decreasing) on (a, b) , then the functions $(f(x) - f(a))/(g(x) - g(a))$ and $(f(x) - f(b))/(g(x) - g(b))$ are also increasing (or decreasing) on (a, b) . If f'/g' is strictly monotone, then the monotonicity in the conclusion is also strict.*

LEMMA 2 ([2, Lemma 4]). *For $x > 0$, the function*

$$k(x) = \frac{\sinh x - x \cosh x}{x^2 \sinh x}$$

is strictly increasing.

Additionally, we prove the following auxiliary results which can be of independent interest.

LEMMA 3 (Sharp upper bound for hyperbolic sinc). *For $x \neq 0$, it is true that*

$$(2.4) \quad \frac{\sinh x}{x} < \sqrt{\frac{(x + \sinh x)(1 + \cosh x)}{4x}}.$$

PROOF. Due to the symmetry of the functions involved at both sides, it suffices to prove (2.4) for $x > 0$. We first consider

$$f(x) = x^2 + x \sinh x - 4 \cosh x + 4.$$

Then

$$f'(x) = 2x + x \cosh x - 3 \sinh x > 0,$$

due to well-known Cusa–Huygens inequality ([2]). Therefore, $f(x)$ is increasing for $x > 0$ and we get $f(x) > f(0)$, i.e.,

$$x^2 + x \sinh x - 4(\cosh x - 1) > 0,$$

which can be written as

$$4(\cosh x - 1)(\cosh x + 1) < x(x + \sinh x)(1 + \cosh x)$$

or

$$4 \sinh^2 x < x(x + \sinh x)(1 + \cosh x).$$

This gives the required inequality (2.4). □

REMARK 1. For $x \neq 0$, it is not difficult to prove

$$\sqrt{\frac{(x + \sinh x)(1 + \cosh x)}{4x}} < \frac{2 + \cosh x}{3}.$$

Thus, we have

$$(2.5) \quad \frac{\sinh x}{x} < \sqrt{\frac{(x + \sinh x)(1 + \cosh x)}{4x}} < \frac{2 + \cosh x}{3}, \quad x \neq 0.$$

REMARK 2. A double inequality analogous to (2.5) also holds in the case of trigonometric functions. It is stated as

$$(2.6) \quad \frac{\sin x}{x} < \sqrt{\frac{(x + \sin x)(1 + \cos x)}{4x}} < \frac{2 + \cos x}{3}, \quad x \neq 0.$$

We skip the proof of (2.6) because it is very similar to that of (2.5).

LEMMA 4 (Refined lower bound for Wilker-type inequality). *For $x \neq 0$, it is true that*

$$(2.7) \quad \left(\frac{x}{\sinh x}\right)^2 + \frac{x}{\tanh x} > 2 + \frac{x(\cosh x - 1)(\sinh x - x)}{2 \sinh^2 x} \\ = 2 + \frac{x(\sinh x - x)}{2(1 + \cosh x)} > 2.$$

PROOF. It is enough to prove that

$$\left(\frac{x}{\sinh x}\right)^2 + \frac{x}{\tanh x} > 2 + \frac{x(\cosh x - 1)(\sinh x - x)}{2 \sinh^2 x}.$$

Equivalently, it corresponds to

$$2x^2 + 2x \sinh x \cosh x > 4 \sinh^2 x + x(\cosh x - 1)(\sinh x - x)$$

or

$$x^2 + x \sinh x \cosh x - 4 \sinh^2 x + x \sinh x + x^2 \cosh x > 0,$$

i.e.,

$$x(x + \sinh x)(1 + \cosh x) > 4 \sinh^2 x,$$

which is true by Lemma 3. □

REMARK 3. The inequality (2.7) is a refinement of the Wilker-type inequality for hyperbolic functions established by Wu and Debnath ([19]).

REMARK 4. It is interesting to see that the circular counterpart of (2.7) is also true for all non-zero real numbers. It is stated as follows:

$$(2.8) \quad \left(\frac{x}{\sin x}\right)^2 + \frac{x}{\tan x} > 2 + \frac{x(\cos x - 1)(\sin x - x)}{2 \sin^2 x} \\ = 2 + \frac{x(x - \sin x)}{2(1 + \cos x)} > 2, \quad x \neq 0.$$

The proof of (2.8) is quite similar to that of (2.7). The importance of (2.7) lies in the fact that it is the sharpest Wilker-type inequality of its kind so far in the literature, and it holds for all non-zero real numbers, although its sharpness can be observed in $(0, \pi)$ and $(-\pi, 0)$ only.

LEMMA 5. *For $x \neq 0$, it is true that*

$$\left(\frac{x}{\sinh x}\right)^2 + \frac{x}{\tanh x} > 2 + \frac{x[\cosh(2x/p) - 1][\sinh(2x/p) - 2x/p]}{p \cdot \sinh^2(2x/p)}$$

if $p \geq 2$.

PROOF. Let

$$g(x) = 2 + x \tanh x - (x \operatorname{sech} x)^2.$$

Differentiation yields

$$\begin{aligned} g'(x) &= \tanh x + 2x^2 \operatorname{sech}^2 x \tanh x - x \operatorname{sech}^2 x \\ &= \frac{x}{\cosh x} \left(\frac{\sinh x}{x} - \frac{1}{\cosh x} \right) + 2x^2 \operatorname{sech}^2 x \tanh x > 0. \end{aligned}$$

Hence, $g(x)$ is increasing and we have that $g(x/2) \geq g(x/p)$ if $x/2 \geq x/p$, i.e., $p \geq 2$. Now, we have

$$\begin{aligned} g\left(\frac{x}{2}\right) &= 2 + \frac{x}{2} \tanh\left(\frac{x}{2}\right) - \left(\frac{x}{2} \operatorname{sech}\left(\frac{x}{2}\right)\right)^2 \\ &= 2 - \frac{x^2}{4 \cosh^2(x/2)} + \frac{x \sinh(x/2)}{2 \cosh(x/2)} \\ &= 2 - \frac{x^2}{2(1 + \cosh x)} + \frac{x \sinh x}{2(1 + \cosh x)} \\ &= 2 + \frac{x(\sinh x - x)}{2(1 + \cosh x)} \\ &= 2 + \frac{x(\cosh x - 1)(\sinh x - x)}{2 \sinh^2 x}. \end{aligned}$$

Similarly, we establish that

$$g\left(\frac{x}{p}\right) = 2 + \frac{x[\cosh(2x/p) - 1][\sinh(2x/p) - 2x/p]}{p \cdot \sinh^2(2x/p)}.$$

By making use of Lemma 4, the conclusion of Lemma 5 follows. \square

3. Main results

We are now in a position to state and prove our main results.

THEOREM 1. For $p > 1$, we define $\phi_p: (0, \pi/2] \rightarrow \mathbb{R}$ by

$$\phi_p(x) = \frac{\ln \left[\frac{\sin x}{x \cos(x/p)} \right]}{x^2}.$$

Then

1. ϕ_p is strictly increasing if $p \leq \sqrt{3}$,
2. ϕ_p is strictly decreasing if $p \geq 2$.

PROOF. We write

$$\phi_p(x) = \frac{\ln[(\sin x)/x] - \ln[\cos(x/p)]}{x^2} = \frac{(\phi_1)_p(x)}{\phi_2(x)},$$

where $(\phi_1)_p(x) = \ln[(\sin x)/x] - \ln[\cos(x/p)]$ and $\phi_2(x) = x^2$, with $(\phi_1)_p(0+) = 0 = \phi_2(0)$. By differentiation, we get

$$\begin{aligned} \frac{(\phi_1)_p'(x)}{\phi_2'(x)} &= \frac{1}{2p} \left[p \frac{x \cos x - \sin x}{x^2 \sin x} + \frac{\tan(x/p)}{x} \right] \\ &= \frac{1}{2p} \left[\frac{\tan(x/p)}{x} + p \frac{\cot x}{x} - \frac{p}{x^2} \right]. \end{aligned}$$

Utilizing (2.1) and (2.2), we obtain

$$\begin{aligned} \frac{(\phi_1)_p'(x)}{\phi_2'(x)} &= \frac{1}{2p} \left[\sum_{n=1}^{\infty} \frac{2^{2n}(2^{2n}-1)}{p^{2n-1} \cdot (2n)!} |B_{2n}| x^{2n-2} - \sum_{n=1}^{\infty} \frac{p \cdot 2^{2n}}{(2n)!} |B_{2n}| x^{2n-2} \right] \\ &= \frac{1}{2p} \sum_{n=1}^{\infty} \frac{2^{2n}}{(2n)!} \left(\frac{2^{2n}-1}{p^{2n-1}} - p \right) |B_{2n}| x^{2n-2}. \end{aligned}$$

By Lemma 1, ϕ_p will be strictly increasing if $(2^{2n}-1)/p^{2n-1} - p > 0$, i.e., $2^{2n}-1 > p^{2n}$ or $p < h(n) := (2^{2n}-1)^{1/2n}$. And it is easy to show that $h(n)$ is strictly increasing for $n = 1, 2, \dots$. This implies that $p \leq \inf\{h(n) : n = 1, 2, \dots\} = h(1) = \sqrt{3}$. Similarly, we can say that ϕ_p will be strictly decreasing if we have $(2^{2n}-1)/p^{2n-1} - p < 0$, or $p > h(n)$. So, we get $p \geq \sup\{h(n) : n = 1, 2, \dots\} = \lim_{n \rightarrow \infty} h(n) = 2$. This completes the proof of Theorem 1. \square

Next, by l'Hôpital's rule, $\phi_p(0+) = \lim_{x \rightarrow 0} \phi_p(x) = 1/(2p^2) - 1/6$, and $\phi_p(\pi/2) = (4/\pi^2) \ln [2/[\pi \cos(\pi/(2p))]]$. Hence, we immediately deduce the following corollaries:

COROLLARY 1. *If $1 < p \leq \sqrt{3}$ and $0 < x \leq \pi/2$, then the best possible constants α_1 and β_1 such that the inequalities*

$$\cos\left(\frac{x}{p}\right) e^{\alpha_1 x^2} < \frac{\sin x}{x} < \cos\left(\frac{x}{p}\right) e^{\beta_1 x^2}$$

hold are $1/(2p^2) - 1/6$ and $(4/\pi^2) \ln [2/[\pi \cos(\pi/(2p))]]$, respectively.

COROLLARY 2. *If $p \geq 2$ and $0 < x \leq \pi/2$, then the inequalities*

$$\cos\left(\frac{x}{p}\right) e^{\beta_1 x^2} < \frac{\sin x}{x} < \cos\left(\frac{x}{p}\right) e^{\alpha_1 x^2}$$

hold with the best possible constants α_1 and β_1 which are as defined in the Corollary 1.

An analogous result involving hyperbolic functions is formulated in the following theorem.

THEOREM 2. *For $p > 0$ and $r > 0$, we define a function $\psi_p: (0, r) \rightarrow \mathbb{R}$ by*

$$\psi_p(x) = \frac{\ln \left[\frac{\sinh x}{x \cosh(x/p)} \right]}{x^2}.$$

Then ψ_p is strictly decreasing if $p \geq 2$. In particular, if $p \geq 2$, then the best possible constants α_2 and β_2 such that the inequalities

$$(3.1) \quad \cosh\left(\frac{x}{p}\right) e^{\alpha_2 x^2} < \frac{\sinh x}{x} < \cosh\left(\frac{x}{p}\right) e^{\beta_2 x^2}, \quad 0 < x < r,$$

hold are $\ln((\sinh r)/[r \cosh(r/p)]) / r^2$ and $1/6 - 1/(2p^2)$, respectively.

PROOF. Set $(\psi_1)_p(x) = \ln[(\sinh x)/x] - \ln[\cosh(x/p)]$ and $\psi_2(x) = x^2$. Clearly $(\psi_1)_p(0+) = 0 = \psi_2(0)$ and $\psi_p(x) = (\psi_1)_p(x)/\psi_2(x)$. In view of using Lemma 1, we differentiate and obtain

$$\frac{(\psi_1)'_p(x)}{\psi'_2(x)} = \frac{1}{2p} \left[p \frac{\coth x}{x} - \frac{p}{x^2} - \frac{\tanh(x/p)}{x} \right] := \frac{1}{2p} (\psi_3)_p(x).$$

Then, we get

$$\begin{aligned}
 (\psi_3)'_p(x) &= -\frac{p}{x} \operatorname{cosech}^2 x - \frac{p}{x^2} \coth x + \frac{2p}{x^3} - \frac{1}{px} \operatorname{sech}^2 \left(\frac{x}{p} \right) + \frac{1}{x^2} \tanh \left(\frac{x}{p} \right) \\
 &= -\frac{p}{x^3} \left[\left(\frac{x}{\sinh x} \right)^2 + \frac{x}{\tanh x} - 2 + \left(\frac{x}{p} \operatorname{sech} \left(\frac{x}{p} \right) \right)^2 - \frac{x}{p} \tanh \left(\frac{x}{p} \right) \right] \\
 &= -\frac{p}{x^3} \left[\left(\frac{x}{\sinh x} \right)^2 + \frac{x}{\tanh x} - 2 \right. \\
 &\quad \left. - \frac{x[\cosh(2x/p) - 1][\sinh(2x/p) - 2x/p]}{p \cdot \sinh^2(2x/p)} \right].
 \end{aligned}$$

By Lemma 1 and Lemma 5, we conclude that ψ_p is strictly decreasing if $p \geq 2$. Consequently,

$$\psi_p(0+) > \psi_p(x) > \psi_p(r-), \quad 0 < x < r.$$

The desired inequalities (3.1) follow due to the limits $\psi_p(0+) = 1/6 - 1/(2p^2)$ and $\psi_p(r-) = \ln[(\sinh r)/[r \cosh(r/p)]]/r^2$. \square

THEOREM 3. For $p > 1$, we define $\varphi_p: (0, \pi/2] \rightarrow \mathbb{R}$ by

$$\varphi_p(x) = \frac{\ln \left[\frac{\sin x}{x \cosh(x/p)} \right]}{x^2}.$$

Then φ_p is strictly decreasing if $p \geq 2$. In particular, if $p \geq 2$, then the best possible constants α_3 and β_3 such that the inequalities

$$(3.2) \quad \cosh \left(\frac{x}{p} \right) e^{\alpha_3 x^2} < \frac{\sin x}{x} < \cosh \left(\frac{x}{p} \right) e^{\beta_3 x^2}, \quad 0 < x \leq \frac{\pi}{2}$$

hold are $(4/\pi^2) \ln [2/[\pi \cosh(\pi/(2p))]]$ and $-[1/(2p^2) + 1/6]$, respectively.

PROOF. We begin with

$$\varphi_p(x) = \frac{\ln(\sin x/x) - \ln(\cosh x/p)}{x^2} = \frac{(\varphi_1)_p(x)}{\varphi_2(x)},$$

where $(\varphi_1)_p(x) = \ln[(\sin x)/x] - \ln[\cosh(x/p)]$ and $\varphi_2(x) = x^2$ with $(\varphi_1)_p(0+) = 0 = \varphi_2(0)$. Differentiation gives

$$\begin{aligned} \frac{(\varphi_1)'_p(x)}{\varphi'_2(x)} &= \frac{1}{2p} \left[p \frac{x \cos x - \sin x}{x^2 \sin x} - \frac{\tanh(x/p)}{x} \right] \\ &= \frac{1}{2p} \left[p \frac{\cot x}{x} - \frac{p}{x^2} - \frac{\tanh(x/p)}{x} \right]. \end{aligned}$$

Utilizing (2.1) and (2.3), we obtain

$$\begin{aligned} \frac{(\varphi_1)'_p(x)}{\varphi'_2(x)} &= \frac{1}{2p} \left[- \sum_{n=1}^{\infty} \frac{p \cdot 2^{2n}}{(2n)!} |B_{2n}| x^{2n-2} - \sum_{n=1}^{\infty} \frac{2^{2n}(2^{2n}-1)}{p^{2n-1} \cdot (2n)!} B_{2n} x^{2n-2} \right] \\ &= -\frac{1}{2p} \sum_{n=1}^{\infty} \frac{2^{2n}}{(2n)!} \left[p \cdot |B_{2n}| - \frac{(2^{2n}-1)}{p^{2n-1}} B_{2n} \right] x^{2n-2} \\ &:= -\frac{1}{2p} \sum_{n=1}^{\infty} \frac{2^{2n}}{(2n)!} a_n x^{2n-2}, \end{aligned}$$

where $a_n = p \cdot |B_{2n}| - [(2^{2n}-1)/p^{2n-1}] B_{2n}$. By Lemma 1, φ_p will be strictly decreasing if $a_n > 0$. But, a_n is always positive for $B_{2n} < 0$ irrespective of p . So we consider the case when $B_{2n} > 0$. In this case, $a_n > 0$ implies that $|B_{2n}| > [(2^{2n}-1)/p^{2n}] B_{2n}$ or $p^{2n} > (2^{2n}-1)$, i.e., $p > (2^{2n}-1)^{1/(2n)} := h(n)$ and $h(n)$ being strictly increasing, we write $p \geq \sup\{h(n) : n = 1, 2, \dots\} = \lim_{n \rightarrow \infty} h(n) = 2$. Finally, $\varphi_p(0+) > \varphi_p(x) > \varphi_p(\pi/2-)$, and the limits $\varphi_p(0+) = -[1/(2p^2) + 1/6]$ and $\varphi_p(\pi/2-) = (4/\pi^2) \ln[2/[\pi \cosh(\pi/(2p))]]$ give the inequalities (3.2). \square

THEOREM 4. For $p \geq 1$ and $r < \pi p/2$, we define $\chi_p : (0, r) \rightarrow \mathbb{R}$ by

$$\chi_p(x) = \frac{\ln \left[\frac{x}{\sinh x \cos(x/p)} \right]}{x^2}.$$

Then χ_p is strictly increasing. In particular, if $p \geq 1$, then the best possible constants α_4 and β_4 such that the inequalities

$$(3.3) \quad \frac{e^{\alpha_4 x^2}}{\cos(x/p)} < \frac{\sinh x}{x} < \frac{e^{\beta_4 x^2}}{\cos(x/p)}, \quad 0 < x < r$$

hold are $-\ln[r/[(\sinh r)(\cos(r/p))]]/r^2$ and $1/6 - 1/(2p^2)$, respectively.

PROOF. We have

$$\chi_p(x) = \frac{(\chi_1)_p(x) - (\chi_1)_p(0+)}{\chi_2(x) - \chi_2(0)},$$

where $(\chi_1)_p(x) = \ln(x/\sinh x) - \ln[\cos(x/p)]$ and $\chi_2(x) = x^2$ with $(\chi_1)_p(0+) = 0 = \chi_2(0)$. Differentiation yields

$$\frac{(\chi_1)'_p(x)}{\chi_2'(x)} = \frac{1}{2} \frac{\sinh x - x \cosh x}{x^2 \sinh x} + \frac{1}{2p^2} \frac{\tan(x/p)}{(x/p)},$$

which is strictly increasing because of Lemma 2 and the fact that $(\tan x)/x$ is strictly increasing in $(0, \pi/2)$. Applying Lemma 1, we conclude that χ_p is strictly increasing in $(0, r)$. Hence, $\chi_p(0+) < \chi_p(x) < \chi_p(r)$ and the desired inequalities (3.3) can be obtained from this and the limits $\chi_p(0+) = 1/(2p^2) - 1/6$ and $\chi_p(r) = \ln[r/[(\sinh r)(\cos(r/p))]]/r^2$. The proof is completed. \square

4. Some particular cases

In this section, we obtain some sharp inequalities from our main results by assigning appropriate values to a parameter p therein. We list the inequalities for sinc and hyperbolic sinc function as follows.

Putting $p = \sqrt{3}$ in Corollary 1 gives

$$(4.1) \quad \cos\left(\frac{x}{\sqrt{3}}\right) < \frac{\sin x}{x} < \cos\left(\frac{x}{\sqrt{3}}\right) e^{\beta_1 x^2}, \quad 0 < x \leq \frac{\pi}{2},$$

where $\beta_1 = (4/\pi^2) \ln[2/(\pi \cos(\pi/2\sqrt{3}))] \approx 0.013219$. This includes the left inequality of (1.1). Putting $p = 2$ in Corollary 2, we obtain

$$(4.2) \quad \cos\left(\frac{x}{2}\right) e^{\beta_1^* x^2} < \frac{\sin x}{x} < \cos\left(\frac{x}{2}\right) e^{-x^2/24}, \quad 0 < x \leq \frac{\pi}{2},$$

where $\beta_1^* = (4/\pi^2) \ln(2\sqrt{2}/\pi) \approx -0.042558$. Lower and upper bounds of (4.2) are sharper than the corresponding lower and upper bounds of (1.1) in the intervals $(\varsigma, \pi/2]$ and $(0, \varsigma)$, respectively, where $\varsigma \approx 1.5204$ and $\varsigma \approx 0.4633$. An upper bound of (4.2) is sharper than that of (4.1) in $(0, \varsigma_1)$, where $\varsigma_1 \approx 1.5346$. The double inequality (4.2) is a complete refinement of (1.3) and (1.6) and it also refines corresponding lower and upper bound of (1.4) in the intervals $(\varsigma_1, \pi/2)$ and $(0, \varsigma_2)$, respectively, where $\varsigma_1 \approx 0.705$ and $\varsigma_2 \approx 1.4372$. Some of




Figure 1. Visual comparison of lower bounds for $(\sin x)/x$ with $x \in [0.8, 1]$; the obtained lower bound is in lightblue color

these facts are illustrated in Figure 1 and Figure 2 for the lower and upper bounds of $(\sin x)/x$, respectively.




Figure 2. Visual comparison of upper bounds for $(\sin x)/x$ with $x \in [0.8, 0.85]$; the obtained upper bound is in light blue color

From Figure 1 and Figure 2, it is clear that the obtained bounds for $(\sinh x)/x$ significantly improve some established bounds of the literature.

Putting $p = 2$ in Theorem 2 yields

$$(4.3) \quad \cosh\left(\frac{x}{2}\right) e^{\alpha_2 x^2} < \frac{\sinh x}{x} < \cosh\left(\frac{x}{2}\right) e^{x^2/24}, \quad 0 < x \leq r,$$

where $\alpha_2 = \ln[(\sinh r)/(r \cosh(r/2))]/r^2$. The inequalities (4.3) uniformly refine (1.7). An upper bound of (4.3) is also a uniform refinement of (1.5). The lower bound of (4.3) is better than the corresponding lower bounds in (1.2) and (1.5) for smaller values of r . However, there is no strict comparison in this case for $(0, r)$. Several other inequalities can be obtained and compared with existing inequalities.

Figure 3 illustrates the sharpness of the obtained upper bound. From Figure 3, we see that the gain of the obtained upper bound in the sharpness sense is consequent.




Figure 3. Visual comparison of lower bounds for $(\sinh x)/x$ with $x \in [0, 3]$; the obtained upper bound is in blue color

NOTE. Due to the symmetry of the functions involved all the inequalities which are true in $(0, \delta)$ are also true in $(-\delta, 0)$.

References

- [1] G.D. Anderson, M.K. Vamanamurthy, and M. Vuorinen, *Conformal Invariants, Inequalities and Quasiconformal Maps*, John Wiley & Sons, New York, 1997.
- [2] Y.J. Bagul and C. Chesneau, *Refined forms of Oppenheim and Cusa-Huygens type inequalities*, Acta Comment. Univ. Tartu. Math. **24** (2020), no. 2, 183–194.
- [3] Y.J. Bagul, R.M. Dhaigude, M. Kostić, and C. Chesneau, *Polynomial-exponential bounds for some trigonometric and hyperbolic functions*, Axioms **10** (2021), no. 4, Paper No. 308, 10 pp.
- [4] B. Chaouchi, V.E. Fedorov, and M. Kostić, *Monotonicity of certain classes of functions related with Cusa-Huygens inequality*, Chelyab. Fiz.-Mat. Zh. **6** (2021), no. 3, 31–337.
- [5] X.-D. Chen, H. Wang, J. Yu, Z. Cheng, and P. Zhu, *New bounds of Sinc function by using a family of exponential functions*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM **116** (2022), no. 1, Paper No. 16, 17 pp.
- [6] C. Chesneau and Y.J. Bagul, *A note on some new bounds for trigonometric functions using infinite products*, Malays. J. Math. Sci. **14** (2020), no. 2, 273–283.
- [7] A.R. Chouikha, C. Chesneau, and Y.J. Bagul, *Some refinements of well-known inequalities involving trigonometric functions*, J. Ramanujan Math. Soc. **36** (2021), no. 3, 193–202.
- [8] I.S. Gradshteyn and I.M. Ryzhik, *Table of Integrals, Series and Products*, Seventh edition, Elsevier/Academic Press, Amsterdam, 2007.
- [9] K.S.K. Iyengar, B.S. Madhava Rao, and T.S. Nanjundiah, *Some trigonometrical inequalities*, Half-Yearly J. Mysore Univ. Sect. B., N.S. **6** (1945), 1–12.
- [10] R. Klén, M. Visuri, and M. Vuorinen, *On Jordan type inequalities for hyperbolic functions*, J. Inequal. Appl. **2010**, Art. ID 362548, 14 pp.
- [11] Y. Lv, G. Wang, and Y. Chu, *A note on Jordan type inequalities for hyperbolic functions*, Appl. Math. Lett. **25** (2012), no. 3, 505–508.
- [12] B. Malešević, T. Lutovac, and B. Banjac, *One method for proving some classes of exponential analytical inequalities*, Filomat **32** (2018), no. 20, 6921–6925.
- [13] B. Malešević and B. Mihailović, *A minimax approximant in the theory of analytic inequalities*, Appl. Anal. Discrete Math. **15** (2021), no. 2, 486–509.
- [14] D.S. Mitrinović, *Analytic Inequalities*, Springer-Verlag, Berlin, 1970.
- [15] E. Neuman and J. Sándor, *On some inequalities involving trigonometric and hyperbolic functions with emphasis on the Cusa-Huygens, Wilker, and Huygens inequalities*, Math. Inequal. Appl. **13** (2010), no. 4, 715–723.
- [16] E. Neuman and J. Sándor, *Inequalities for hyperbolic functions*, Appl. Math. Comput. **218** (2012), no. 18, 9291–9295.
- [17] C. Qian, X.-D. Chen, and B. Malešević, *Tighter bounds for the inequalities of Sinc function based on reparameterization*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM **116** (2022), no. 1, Paper No. 29, 38 pp.
- [18] J. Sándor, *Two applications of the Hadamard integral inequality*, Notes Number Theory Discrete Math. **23** (2017), no. 4, 52–55.
- [19] S. Wu and L. Debnath, *Wilker-type inequalities for hyperbolic functions*, Appl. Math. Lett. **25** (2012), no. 5, 837–842.
- [20] Z.-H. Yang, *New sharp bounds for logarithmic mean and identric mean*, J. Inequal. Appl. **2013**, 2013:116, 17 pp.
- [21] Z.-H. Yang, *Refinements of a two-sided inequality for trigonometric functions*, J. Math. Inequal. **7** (2013), no. 4, 601–615.
- [22] Z.-H. Yang and Y.-M. Chu, *Jordan type inequalities for hyperbolic functions and their applications*, J. Funct. Spaces **2015**, Art. ID 370979, 4 pp.
- [23] L. Zhang and X. Ma, *Some new results of Mitrinović-Cusa’s and related inequalities based on the interpolation and approximation method*, J. Math. **2021**, Art. ID 5595650, 13 pp.

- [24] L. Zhu, *Generalized Lazarevic's inequality and its applications-Part II*, J. Inequal. Appl. **2009**, Art. ID 379142, 4 pp.
- [25] L. Zhu, *Some new bounds for Sinc function by simultaneous approximation of the base and exponential functions*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **114** (2020), no. 2, Paper No. 81, 17 pp.
- [26] L. Zhu and R. Zhang, *New inequalities of Mitrinović-Adamović type*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Mat. RACSAM **116** (2022), no. 1, Paper No. 34, 15 pp.

YOGESH J. BAGUL
DEPARTMENT OF MATHEMATICS
K. K. M. COLLEGE, MANWATH
DIST: PARBHANI(M.S.)-431505
INDIA
e-mail: yjbagul@gmail.com

SUMEDH B. THOOL
DEPARTMENT OF MATHEMATICS
GOVERNMENT VIDARBHA INSTITUTE OF SCIENCE AND HUMANITIES
AMRAVATI(M. S.)-444604
INDIA
e-mail: sumedhmaths@gmail.com

CHRISTOPHE CHESNEAU
LMNO
UNIVERSITY OF CAEN-NORMANDIE
CAEN
FRANCE
e-mail: christophe.chesneau@unicaen.fr

RAMKRISHNA M. DHAIGUDE
DEPARTMENT OF MATHEMATICS
GOVERNMENT VIDARBHA INSTITUTE OF SCIENCE AND HUMANITIES
AMRAVATI(M. S.)-444604
INDIA
e-mail: rmdhaigude@gmail.com